

1. (a) Given that  $S^n$  is defined as the boundary of the closed unit ball in  $\mathbb{R}^{n+1}$ , describe  $S^0$ .
- (b) The intersection of  $S^2$  with the  $xy$ -plane in  $\mathbb{R}^3$  is  $S^1$ , a circle of radius 1. We call this a *great circle*, since no other circle on  $S^2$  has a larger radius. Similarly, the intersection of  $S^2$  with any plane through the origin in  $\mathbb{R}^3$  is called a **great circle**. What is the intersection of two great circles?
- (c) Now one dimension higher. Denote points in  $\mathbb{R}^4$  by  $(x, y, z, w)$ . Prove *rigorously* that the intersection of  $S^3$  with the  $xyz$ -hyperplane in  $\mathbb{R}^4$  is  $S^2$ . We call such a 2-sphere a **great sphere**.
- (d) What is the intersection of two great spheres? Prove your answer rigorously for the intersection of the great sphere cut out by the  $xyz$ -hyperplane with the great sphere cut out by the  $yzw$ -hyperplane.
- (e) The general form equation of a plane in  $\mathbb{R}^3$  is:  $ax + by + cz = d$ , where  $a, b, c, d$  are constants. The general form equation of an  $n - 1$ -dimensional hyperplane in  $\mathbb{R}^n$  is:  $a_1x_1 + a_nx_n = b$ , where  $a_i$  and  $b$  are constants. Think this way: in  $\mathbb{R}^n$ , you have  $n$  “degrees of freedom”. An equation gives one constraint, reducing the number of degrees of freedom to  $n - 1$ ; hence the solutions to the equation form an  $(n - 1)$ -dimensional manifold (which is a hyperplane if the equation is linear). Now, here’s the question: What is the intersection of two  $(n - 1)$ -dimensional hyperplanes that pass through the origin in  $\mathbb{R}^n$ ? Explain your reasoning.
- (f) Give a definition for a great  $(n - 1)$ -sphere in  $S^n \subset \mathbb{R}^{n+1}$ . Describe the intersection of two great  $(n - 1)$ -spheres in  $S^n$ , and explain your reasoning (it doesn’t have to be a rigorous proof, but only a clear and convincing explanation; though a rigorous proof wouldn’t be bad either!). Hint: Use the previous part of this problem.
2. A torus  $T^2$  can be defined as a “solid square”  $I^2 = [0, 1] \times [0, 1]$  with its opposite edges identified (with the “appropriate” orientations):  $T^2 = I^2/R$ , where  $R$  denotes the equivalence relation  $(x, 0) \sim (x, 1)$ ,  $(0, y) \sim (1, y)$ . Similarly, a 3-dimensional torus  $T^3$  can be defined as a solid cube  $I^3$  with its opposite faces identified (with the “appropriate” orientations). Make this precise by giving an appropriate definition for  $R'$ :  $T^3 = I^3/R'$ , where  $R'$  is  $\dots$ .
3.  $T^2$  can also be defined as  $S^1 \times S^1$ . We can informally explain how this definition is equivalent to the above definition ( $T^2 = I^2/R$ ) by arguing as follows. For every  $t \in I$ , the two endpoints of  $I \times \{t\} \subset I^2$  are identified into one point; so each  $I \times \{t\}$  becomes homeomorphic to  $S^1 \times \{t\}$ . Therefore,  $I^2/R$  is homeomorphic to  $S^1 \times I$  with  $S^1 \times \{0\}$  identified with  $S^1 \times \{1\}$  (with the “right” orientation). Thus, we get  $S^1 \times S^1$ . Give a similar informal argument to show  $I^3/R' \simeq S^1 \times S^1 \times S^1$ .
4. (a) What familiar space is a punctured 3-sphere ( $S^3$  minus one point) homeomorphic to? Briefly explain why.
- (b) Let  $p$  be an arbitrary point in  $S^2$ . Then, in  $S^2 \times S^1$ ,  $\{p\} \times S^1$  is a simple closed curve. Draw a schematic diagram of this. Call this simple closed curve  $C$ . Does  $C$  bound a disk in  $S^2 \times S^1$ ?
- (c) What familiar space is  $(S^2 \times S^1) - (N_\epsilon(C))^\circ$  (i.e.,  $(S^2 \times S^1)$  minus the interior of an  $\epsilon$ -neighborhood of  $C$ ) homeomorphic to?
5. (a) It is possible to travel in  $\mathbb{R}^3$  from the point  $(-1, 0, 0)$  to the point  $(1, 0, 0)$  by walking along straight line segments and without ever touching the  $y$ -axis. Explain how. How about without ever the  $yz$ -plane?
- (b) It is possible to travel in  $\mathbb{R}^4$  from the point  $(-1, 0, 0, 0)$  to the point  $(1, 0, 0, 0)$  by walking along straight line segments and without ever touching the  $yz$ -plane  $\{(x, y, z, w) \in \mathbb{R}^4 : x = w = 0\}$ . Explain how.

6. A **2-component link** consists of two disjoint circles embedded in  $\mathbb{R}^3$ . For example, let  $X \subset \mathbb{R}^3$  be the unit circle in the  $xy$ -plane centered at the origin, and let  $Y \subset \mathbb{R}^3$  be the unit circle in the  $yz$ -plane centered at  $(0, 1, 0)$ . Then  $X \cup Y$  is a 2-component link; in fact, it has its own name: the **Hopf link** (named after a mathematician). Draw a picture of this link, showing the three axes in  $\mathbb{R}^3$  and the coordinates of the circles' centers. The two components of the link,  $X$  and  $Y$ , cannot be “pulled apart”; more precisely, if we let  $Y'$  be a unit circle centered at  $(0, 5, 0)$ , then  $X \cup Y$  is not isotopic to  $X \cup Y'$ . This is not easy to prove rigorously with what we know so far, but should be clear intuitively—do you see it?

Now,  $\mathbb{R}^3$  can be viewed as a subset of  $\mathbb{R}^4$ :  $\mathbb{R}^3 = \{(x, y, z, w) \in \mathbb{R}^4 : w = 0\}$ . Then  $X \cup Y \subset \mathbb{R}^3 \subset \mathbb{R}^4$ . Explain informally how  $X \cup Y$  can be “pulled apart” in  $\mathbb{R}^4$ .

7. Give a detailed and rigorous proof that if  $n > m$ , then  $S^n$  cannot be embedded in  $\mathbb{R}^m$ . (Use the theorem that says  $S^n$  cannot be embedded in  $\mathbb{R}^n$ .)
8. Prove or disprove: If two connected topological spaces are glued together, the result is connected. Here's a precise statement: Let  $X$  and  $Y$  be connected topological spaces. Let  $\sim$  be an equivalence relation on  $X \cup Y$  such that for at least one point  $x \in X$  and at least one point  $y \in Y$ ,  $x \sim y$ ; then  $(X \cup Y)/\sim$  is connected.

### Extra Credit Problems

9. (a) Explain how  $S^2 \times S^1$  can be viewed as two solid tori glued together along their boundaries.  
 (b) Explain how  $S^3$  can be viewed as two solid tori glued together along their boundaries. Hint: view it as the boundary of  $B^2 \times B^2$  ( $\simeq B^4$ ).  
 (c) Explain why the above does *not* imply  $S^3 \simeq S^2 \times S^1$ .
10. Give an embedding of  $\mathbb{RP}^2$  into  $\mathbb{R}^4$ .