

Definition 1. A **metric space** M consists of a set X and a **distance function** $d : X \times X \rightarrow [0, \infty)$ such that $\forall x, y, z \in X$

1. $d(x, y) = 0$ iff $x = y$;
2. $d(x, y) = d(y, x)$ (d is symmetric);
3. $d(x, z) \leq d(x, y) + d(y, z)$ (triangle inequality).

Example 1. \mathbb{R} with the **Euclidean metric** (the “standard” metric):

$X = \mathbb{R}$, $d(x, y) = |x - y|$. Why is this a metric space?

If instead we had $d(x, y) = x - y$, would we still have a metric space?

Example 2. \mathbb{R} with the **discrete metric**, denoted \mathbb{R}_d :

$X = \mathbb{R}$, $d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$. Why is this a metric space?

What if we let $d(x, y) = 0$ for all x, y , is it still a metric space?

Example 3. \mathbb{R}^n with the **Euclidean metric** :

$X = \mathbb{R} \times \cdots \times \mathbb{R}$ (n times); for $x = (x_1, \cdots, x_n)$, $y = (y_1, \cdots, y_n)$, $d(x, y) = \sqrt{(x_1 - y_1)^2 + \cdots + (x_n - y_n)^2}$.

Why is this a metric space? Conditions 1 and 2 of the definition (above) are clearly satisfied. Condition 3 is the well-known triangle inequality (skip proof).

Example 4. \mathbb{R}^2 with the **taxicab metric** :

$X = \mathbb{R}^2$, for $a = (a_1, a_2)$, $b = (b_1, b_2)$, $d(a, b) = |a_1 - b_1| + |a_2 - b_2|$. Why is this a metric space? (HW)

Note.

1. Unless stated otherwise, whenever we refer to \mathbb{R} as a metric space without stating what the distance function d is, we mean “ \mathbb{R} with the Euclidean metric.”
2. For a metric space $M = (X, d)$, X is called **the underlying set**. We will often abuse notation and write M instead of X , or vice versa; for example, we may write $x \in M$ instead of $x \in X$; or we may refer to X as a metric space, when it’s really $M = (X, d)$ that’s a metric space.

Definition 2. Given a metric space M , a point $x \in M$, and a real number $r > 0$, the **ball** of radius r around x is defined as

$$B_r(x) = \{y \in M \mid d(x, y) < r\}$$

Example 5. In \mathbb{R} with the Euclidean metric, $B_2(1) = ?$ ¹

Example 6. In \mathbb{R}^2 with the Euclidean metric, what does $B_2(1, 2)$ look like? (Strictly speaking, we should write $B_2((1, 2))$; but too many parentheses can make it difficult to read, so we slightly abuse notation and write only one set of parentheses.) How about $B_2(1, 2) \subset \mathbb{R}^3$, what does it look like?

Example 7. In \mathbb{R}_d , what is $B_3(8)$? What is $B_{0.5}(8)$?²

Example 8. In \mathbb{R}^2 with the taxicab metric, what does $B_1(0, 0)$ look like?

Example 9. Is there a metric on \mathbb{R}^2 for which $B_1(0, 0) = (-1, 1) \times (-1, 1)$?³

Definition 3. A subset A of a metric space M is said to be **open** in M iff $\forall x \in A$, $\exists r > 0$ such that $B_r(x) \subset A$.

¹The open interval from -1 to 3 : $(-1, 3)$.

² $B_3(8) = \mathbb{R}$; $B_{0.5}(8) = \{8\}$.

³ $d(a, b) = \max\{|a_1 - b_1|, |a_2 - b_2|\}$.

Example 10. The interval $(-1, 1]$ is not open in \mathbb{R} . Why? ⁴

Example 11. The interval $(-1, 1)$ is an open subset of \mathbb{R} . Why?

Proof: Given an arbitrary $x \in (-1, 1)$, let $r = \min\{d(x, 1), d(x, -1)\}$. Then, we prove as follows that $B_r(x) \subset (-1, 1)$. Let $y \in B_r(x)$; we'll show $y \in (-1, 1)$. We will do so by showing that $d(0, y) < 1$. By definition of $B_r(x)$, $d(x, y) < r$; so $d(x, y) < \min\{d(x, 1), d(x, -1)\}$; so $d(x, y) < d(x, 1)$ and $d(x, y) < d(x, -1)$. By the triangle inequality, $d(0, y) \leq d(0, x) + d(x, y)$. So, $d(0, y) < d(0, x) + d(x, 1)$ and $d(0, y) < d(0, x) + d(x, -1)$. If $x \geq 0$, then the right hand side of the first inequality equals 1. If $x < 0$, then the left hand side of the second inequality equals 1. So either way, $d(0, y) < 1$, as desired. We showed that for every $x \in (-1, 1)$, there is a positive r such that $B_r(x) \subset (-1, 1)$. So by the definition of open, $(-1, 1)$ is an open subset of \mathbb{R} .

Example 12. Is the interval $(2, \infty)$ open in \mathbb{R} ? Yes. Why?

Definition 4. Let A be a subset of a metric space M . The **complement** of A is $A^c = M - A$. A is said to be **closed** in M iff its complement A^c is open in M .

Example 13. $(-\infty, -1] \cup [1, \infty)$ is closed in \mathbb{R} . Why?

Example 14. Is $(-\infty, -1]$ closed in \mathbb{R} ? ⁵

Example 15. Is $[-1, 1]$ closed in \mathbb{R} ? ⁶

Example 16. $[-1, 1)$ is neither open nor closed in \mathbb{R} . Why?

Example 17. \mathbb{R} is open in \mathbb{R} . Why? ϕ is open in \mathbb{R} . Why?

Example 18. \mathbb{R} is closed in \mathbb{R} . ϕ is closed in \mathbb{R} . Why?

Example 19. Is the x -axis open or closed or neither in \mathbb{R}^2 ? ⁷

Example 20. Find an open set in \mathbb{R}_d . Find a closed set in \mathbb{R}_d . ⁸

(Quote from Munkres's book, *Topology*: Q: "What's the difference between a door and a set?" A: "A door is always either open or closed.")

For emphasis, $B_r(x)$ is sometimes called the *open ball* of radius r around x . In contrast, we have:

Definition 5. The **closed ball** of radius r around x is defined as

$$\overline{B_r(x)} = \{y \in M \mid d(x, y) \leq r\}$$

Example 21. Draw the open and closed balls of radius 5 around the point 2 in \mathbb{R} . Draw the open and closed balls of radius 5 around the point $(2, 5)$ in \mathbb{R}^2 .

Definition 6. Let A be a subset of a metric space M . A point $x \in M$ is said to be a **limit point** of A iff every ball around x contains a point of A other than x .

(Synonyms of *limit point*: cluster point; accumulation point.)

Example 22. Let $M = \mathbb{R}$, $A = [0, 2)$. Which of the points $x = 0, 1, 2, 3$ are limit points of A ? Why? ⁹
What if $A = [0, 1] \cup \{2\}$? ¹⁰

(Equivalent definition of limit point: x is a limit point of A iff $\forall \epsilon > 0, \exists y \in A - \{x\}$ such that $d(x, y) < \epsilon$.)

⁴Because there is no positive r for which $B_r(1) \subset (-1, 1]$.

⁵Yes. Why?

⁶Yes. Why?

⁷Closed. Why?

⁸Each of \mathbb{R}_d and ϕ is both open and closed.

⁹0, 1 and 2.

¹⁰0 and 1.

Theorem 1. A subset A of a metric space M is closed iff it contains all its limit points.

Proof. “ \Rightarrow ” : Suppose A is closed. Then, by definition, A^c is open. Let x be a limit point of A . We want to show $x \in A$. By definition of limit point, every open ball around x intersects $A - \{x\}$; therefore no open ball around x is entirely contained in A^c . This implies $x \notin A^c$, since if x were in A^c , then there would be an open ball around x contained entirely in A^c (since A^c is open). Finally, since $x \notin A^c$, x must be in A , as desired.

“ \Leftarrow ” : (Do yourself!) □

Definition 7. Given a subset A of a metric space M , its **interior** A° is defined as the set of all points $x \in A$ such that some open ball around x is a subset of A . (A° is also written as $\text{Int } A$ or $\text{int}(A)$.)

Example 23. (a) What is the interior of $[2, 5) \subset \mathbb{R}$? ¹¹

(b) What is the interior of $(2, 5) \subset \mathbb{R}$? ¹²

(c) What is the interior of the closed ball of radius 2 around the origin in \mathbb{R}^2 ? ¹³

Definition 8. Given a subset A of a metric space M , its **closure** \bar{A} is defined as A union the set of all limit points of A . The **boundary** of A is defined as $\partial A = \bar{A} - A^\circ$.

Example 24. (a) What are the closure and boundary of $[2, 5) \subset \mathbb{R}$? ¹⁴

(b) What is the closure and boundary of the closed ball of radius 2 around the origin in \mathbb{R}^2 ? ¹⁵

Continuity

Definition 9. Let M_1, M_2 be metric spaces, with d_1 and d_2 as their corresponding distance functions. A function $f : M_1 \rightarrow M_2$ is said to be **continuous at** $a \in M_1$ iff as $x \rightarrow a$, $f(x) \rightarrow f(a)$; this means: $\forall \epsilon > 0, \exists \delta > 0$ such that for every x that satisfies $d_1(a, x) < \delta$ we have $d_2(f(a), f(x)) < \epsilon$. We say f is **continuous** if it is continuous at every point in M_1 .

Example 25. Prove that $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 2x$ is continuous.

Proof: Fix an arbitrary point $a \in \mathbb{R}$. We will show f is continuous at a by showing that $\forall \epsilon > 0, \exists \delta > 0$ such that for every x that satisfies $d(a, x) < \delta$ we have $d(f(a), f(x)) < \epsilon$.

Pick any $\epsilon > 0$. Let $\delta = \epsilon/2$. Then, for every x that satisfies $d(a, x) < \delta$ we have: $|a - x| < \delta$, so $|2a - 2x| < 2\delta$, so $d(f(a), f(x)) < \epsilon$, as desired. Since a was an arbitrary point, f is continuous at every point in \mathbb{R} .

Example 26. Determine whether each of the following functions f and g from \mathbb{R} to \mathbb{R} is continuous at 0. (Support your answers informally, without rigorous proof.)

$$f(x) = \begin{cases} \sin(1/x) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases} \quad g(x) = \begin{cases} x \sin(1/x) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

¹¹ $(2, 5)$.

¹² $(2, 5)$.

¹³the open ball of radius 2 around the origin.

¹⁴closure = $[2, 5]$; boundary = $\{2, 5\}$.

¹⁵closure = itself; boundary = circle of radius 2 around the origin.