

Review defs of neighborhood, locally homeomorphic, and manifold.

Recall that, in the definition of manifold, we can replace “locally homeomorphic to an open ball in  $\mathbb{R}^n$ ” with “locally homeomorphic to  $\mathbb{R}^n$ .”

*Example 1.* Is the open rectangle  $(0, 1) \times (0, 2) \subset \mathbb{R}^2$  a manifold? Yes. Of what dimension? 2.

Is the closed rectangle  $[0, 1] \times [0, 2] \subset \mathbb{R}^2$  a manifold? No. Why?

We’d like to say that the closed rectangle is a manifold *with boundary*. Before defining this, we need another definition.

*Definition 1.* The  **$n$ -dimensional upper half-space** is defined as

$$\mathbb{R}_+^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n \geq 0\}$$

When  $n = 2$ ,  $\mathbb{R}_+^2$  is also called the **upper half-plane**.

*Example 2.* Draw a picture of what each of  $\mathbb{R}_+^1$  and  $\mathbb{R}_+^2$  looks like.

*Example 3.* Let  $X = B_1(0, 0) \cap \mathbb{R}_+^2$ . Is  $X$  homeomorphic to  $\mathbb{R}_+^2$ ? Yes. Why? Does every point in  $X$  have a nbhd that’s homeomorphic to either  $\mathbb{R}^2$  or  $\mathbb{R}_+^2$ ? Yes. Why?

*Definition 2.* (Overrides previous definition of manifold) A topological space  $X$  is called an  **$n$ -dimensional manifold** ( $n$ -mfd for short) if it is Hausdorff, Second Countable, and every point  $x \in X$  has a neighborhood that is homeomorphic to  $\mathbb{R}^n$  or  $\mathbb{R}_+^n$ . A point that has a neighborhood homeomorphic to  $\mathbb{R}_+^n$  but not to  $\mathbb{R}^n$  is called a **boundary point**. The set of all such points (if any) is called the **boundary** of  $X$ , denoted by  $\partial X$ . If  $\partial X \neq \emptyset$ , then, for emphasis,  $X$  is sometimes called a **manifold with boundary**.

*Example 4.* Let  $X = ([0, 1] \times [0, 1]) / \{(0, y) \sim (1, y)\}$ . Draw a picture of  $X$ . Is  $X$  a manifold? Yes. Is it a manifold with boundary? Yes. What is  $\partial X$ ? It’s the disjoint union of two circles:  $([0, 1] \times \{0\}) / \{(0, 0) \sim (1, 0)\} \cup ([0, 1] \times \{1\}) / \{(0, 1) \sim (1, 1)\}$ .

*Example 5.* Let  $X = [0, 1] \times [0, 1] / \{(0, y) \sim (1/2, y)\}$ . Draw a picture of  $X$ . Is  $X$  a manifold? No. Why?

*Remark.* Depending on the context, the term boundary can mean two similar but *different* things: when applied to a subset  $A$  of a topological space, it means  $\overline{A} - A^\circ$ ; but when applied to a manifold, it is defined according to the above definition.

*Theorem 1.* (Classification of 1-manifolds) Every 1-manifold is homeomorphic to  $[0, 1]$  or  $(0, 1)$  or  $[0, 1]$  or  $S^1$ .

Idea of Proof: What things can you create by joining or overlapping line segments end-to-end? Only these!

We are often interested in studying manifolds that are compact and have no boundary. (Why? One reason is that non-compact manifolds are usually more difficult to study, or at least different from compact ones.) There is a name for these manifolds:

*Definition 3.* A manifold is said to be **closed** if it is compact and has no boundary.

*Note.* Do not confuse the two different meanings of *closed*; they depend on the context: A subset  $A$  of a topological space  $X$  is closed if  $X - A$  is open in  $X$  (i.e., is in  $\mathcal{T}$ ). A manifold is closed if it’s compact and has no boundary.

*Example 6.* Which of the four 1-mfds are closed? Ans: Only  $S^1$ . Why is each of the other three not closed?

---

### Hausdorff Spaces

Definition: A topological space  $X$  is said to be **Hausdorff** iff every pair of distinct points  $x_1, x_2 \in X$  can be **separated** by open sets, i.e., there exist disjoint open sets  $U_1, U_2 \subset X$  such that  $x_i \in U_i$ .

---

*Example 7.* Q: Is  $\mathbb{R}^2$  with the Euclidean metric Hausdorff? Yes. Why?

Q: Is  $\mathbb{R}^2$  with the discrete topology Hausdorff? Yes. Why?

Q: Is  $\mathbb{R}^2$  with the indiscrete topology Hausdorff? No. Why?

---

*Example 8.* Which of the following are manifolds? Why?

(a)  $([0, 2] \cup [5, 7]) / \{ \forall x \in [0, 1], x \sim (x + 5) \}$ . Ans: Not a mfd, since the point  $\{1, 6\}$  in the quotient space does not have a nbhd that's homeomorphic to  $\mathbb{R}^n$  or  $\mathbb{R}_+^n$  for any  $n$ .

(b)  $([0, 2] \cup [5, 7]) / \{ \forall x \in [0, 1), x \sim (x + 5) \}$ . Ans: Every point does have a nbhd that's homeomorphic to  $\mathbb{R}$ ; nevertheless, this is not a mfd! Why? Because it's not Hausdorff: The points 1 and 6 cannot be separated by open sets.

---