

---

# Differential Equations

Math 341 Fall 2013  
©2013 Ron Buckmire

MWF 12:50-1:45pm Fowler 307  
<http://faculty.oxy.edu/ron/math/341/13/>

---

## Worksheet 16

**TITLE** Straight Line Solutions

**CURRENT READING** Blanchard, 3.2

---

---

**Homework Assignments due Monday October 28** (\* indicates EXTRA CREDIT)

**Section 3.1:** 6, 7, 8, 10, 13, 18\*.

**Section 3.2:** 8, 9, 12, 16\*, 17, 18\*.

---

---

### SUMMARY

Eigenvalues and eigenvectors return from Linear Algebra and are important in the case where Linear Systems of ODEs have solutions that look like straight lines.

---

#### 1. The Significance of Eigenvectors and Eigenvalues

Recall the solutions  $\vec{x}_1(t) = \begin{bmatrix} e^{2t} \\ 0 \end{bmatrix}$  and  $\vec{x}_2(t) = \begin{bmatrix} -e^{-4t} \\ 2e^{-4t} \end{bmatrix}$  to the ODE  $\frac{d\vec{x}}{dt} = \begin{bmatrix} 2 & 3 \\ 0 & -4 \end{bmatrix} \vec{x}$  from *Worksheet #15*.

Notice that  $\vec{x}_1(t) = \begin{bmatrix} e^{2t} \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^{2t}$  and  $\vec{x}_2(t) = \begin{bmatrix} -e^{-4t} \\ 2e^{-4t} \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix} e^{-4t}$ .


#### Question

Do you notice anything interesting about the vectors  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$ ? Any relationship to the matrix  $A = \begin{bmatrix} 2 & 3 \\ 0 & -4 \end{bmatrix}$ ? What happens if you multiply each vector by  $A$ ?

#### Answer

The vectors in question are \_\_\_\_\_.

Consider the direction field for the ODE  $\frac{d\vec{x}}{dt} = \begin{bmatrix} 2 & 3 \\ 0 & -4 \end{bmatrix} \vec{x}$ :



It turns out that the general solution to  $\frac{d\vec{x}}{dt} = \begin{bmatrix} 2 & 3 \\ 0 & -4 \end{bmatrix} \vec{x}$  can be written as

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) = c_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^{2t} + c_2 \begin{bmatrix} -1 \\ 2 \end{bmatrix} e^{-4t}.$$

**Exercise**

On the above direction field, we want to draw in the solutions  $\vec{x}_1(t)$  and  $\vec{x}_2(t)$ . Does it matter what your initial condition is?

What happens as  $t \rightarrow \infty$ ? What about as  $t \rightarrow -\infty$  (i.e. reverse direction of the arrows)? Does one of the solutions seem more “attractive” than the other?

**EXAMPLE**

Consider the system  $\frac{d\vec{x}}{dt} = \begin{bmatrix} 1 & 3 \\ 5 & 3 \end{bmatrix} \vec{x}$ . Find the eigenvalues  $\lambda$  and eigenvectors  $\vec{v}$  of  $\begin{bmatrix} 1 & 3 \\ 5 & 3 \end{bmatrix}$ .

Show that the general solution can be written as  $\vec{x} = c_1\vec{v}_1e^{\lambda_1t} + c_2\vec{v}_2e^{\lambda_2t}$  and confirm that it is actually a solution of  $\vec{x}' = \begin{bmatrix} 1 & 3 \\ 5 & 3 \end{bmatrix} \vec{x}$ .

## 2. General Solution To Homogeneous Linear Systems

**THEOREM**

The general solution  $\vec{x}(t)$  on the interval  $(-\infty, \infty)$  to a homogeneous system of linear DEs  $\frac{d\vec{x}(t)}{dt} = A(t)\vec{x}(t)$  can be written as  $\vec{x} = c_1\vec{v}_1e^{\lambda_1t} + c_2\vec{v}_2e^{\lambda_2t} + c_3\vec{v}_3e^{\lambda_3t} + \dots + c_n\vec{v}_ne^{\lambda_nt}$  where  $\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_n$  and  $\vec{v}_1, \vec{v}_2, \vec{v}_3, \dots, \vec{v}_n$  are the eigenvalues and corresponding eigenvectors of the matrix  $A$ .

### 3. Phase Portraits With Straight Line Solutions

**Exercise**

Solve  $\frac{dx}{dt} = 2x + 2y$ ,  $\frac{dy}{dt} = x + 3y$ .

**GroupWork**

Use `HPGSystemSolver` (or `PPLANE`) to sketch the phase portrait of the linear system

$\frac{d\vec{x}}{dt} = \begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix} \vec{x}$  you solved above, in the space below.