
Numerical Analysis

Math 370 Fall 2002

©2002 Ron Buckmire

MWF 9:30am - 10:25pm

Fowler 127

Worksheet 4: Friday September 13

SUMMARY Taylor Series Approximations and Order of Convergence for Functions

Order of Convergence of Functions

If we know that $\lim_{h \rightarrow 0} F(h) = L$ and $\lim_{h \rightarrow 0} G(h) = 0$ and if a positive constant K exists with

$$|F(h) - L| \leq KG(h), \quad \text{for sufficiently small } h$$

then we write $F(h) = L + \mathcal{O}(G(h))$

This can also be computed using the idea that $F(h) = L + \mathcal{O}(G(h))$ if and only if

$$\lim_{h \rightarrow 0} \frac{|F(h) - L|}{|G(h)|} = K$$

where K is some positive, finite constant.

Example

Show that the expression $\cos(h) + \frac{h^2}{2}$ is $1 + \mathcal{O}(h^4)$

What is the rate of convergence of $\sin(h^3)$ as $h \rightarrow 0$

Taylor Expansions

Another approach to figuring out order of convergence of functions is to use *Taylor Series Approximations* to assist you in satisfying the inequality version of the definition.

Recall that if you have a function $f(x)$ near a point $x = a$ and $f(x)$ is infinitely-differentiable, you can write down

$$f(x) =$$

or you can truncate this series and write down

$$f(x) \approx$$

Suppose we approximate the function $f(x)$ near the point $a = 0$ then we obtain what is known as a *Maclaurin Series*.

Maclaurin Series

$$f(x) = \sum_{k=0}^{\infty} f^{(k)}(0) \frac{x^k}{k!}$$

If we are only interested in the behavior of the function for small values of x near 0, i.e. for $|h| \ll 1$ then we can write the expression as

$$f(0 + h) = \sum_{k=0}^{\infty} f^{(k)}(0) \frac{h^k}{k!}$$

Interestingly, we can write an exact expression for the truncated form of this expression as

$$f(0 + h) = f(0) + f'(0)h + f''(0)\frac{h^2}{2} + \mathcal{O}(h^3)$$

Yes, this last term is the same “big oh” that we have been discussing in regards to order of convergence of a function to its limit.

Exercise

Write down the following Taylor Series Approximations (for small h) for the following functions:

$$\sin(h) \approx$$

$$\cos(h) \approx$$

$$e^h \approx$$

$$(1 + h)^p \approx$$

$$\ln(1 + h) \approx$$

Example

Show that you can use a truncated Maclaurin Expansion to prove that $\cos(h) + \frac{h^2}{2} = 1 + \mathcal{O}(h^4)$

The Three Ways Of Computing Order of Convergence of a Function are

1. **Limit Method**

2. **Bounding/Inequality Method**

3. **Truncated Taylor/Maclaurin Expansion**

(next to each of the methods above, write a short note to yourself explaining the method.)

GROUPWORK

Work in Groups of 2 or 3 to find the limit and express the order of convergence in terms of $f(h) = c + \mathcal{O}(h^\alpha) = c + o(h^\beta)$ for the following:

1. $f(h) = \frac{1 + h - e^h}{h^2}$

2. $f(h) = \frac{1}{1 - h^4}$