

Abstract Submitted  
for the DPP05 Meeting of  
The American Physical Society

Sorting Category: 1.3.5 (Computational)

**Particle Dynamics in Asymmetry-Induced Transport:  
a Computational Study** D.L. EGGLESTON, Occidental College —

We have developed a simple computer code as an aid to resolving the discrepancies between our experiments<sup>1</sup> and the theory<sup>2</sup> of asymmetry-induced transport. The code employs the fourth-order Runge-Kutta method to advance the particles in prescribed fields matching our experiment. For a single helical asymmetry  $\phi(r) \cos(kz + l\theta - \omega t)$ , significant motion in the radial direction is restricted to those particles near the resonant velocity. Both the location and the width of this resonance are consistent with expectations. When a standing wave asymmetry is used (i.e., two counter-propagating helical waves), additional dynamical behaviors are observed. Stochastic motion occurs when the resonant regions of the two waves overlap, allowing a larger population of particles to undergo large radial excursions. There is also a class of particles with restricted axial motion, as in trapped particle modes<sup>3</sup>. These particles, which also make large radial excursions, are located near the radius where  $\dot{\theta} = \omega/l$ . Further progress in understanding asymmetry-induced transport may require inclusion of these effects.

<sup>1</sup>D.L. Eggleston and B. Carrillo, Phys. Plasmas **10**, 1308 (2003).

<sup>2</sup>D.L. Eggleston and T.M. O'Neil, Phys. Plasmas **6**, 2699 (1999).

<sup>3</sup>A.A. Kabantsev et al., Phys. Rev. Lett. **89**, 245001 (2002).

☐  
☒

Prefer Oral Session

Prefer Poster Session

Dennis L. Eggleston  
dleggles@oxy.edu  
Occidental College

Date submitted: July 17, 2006

Electronic form version 1.4