

We previously derived the initial value problem for the motion of a spring.

$$\begin{aligned}x' &= v & x(0) &= a \\v' &= -b^2x & v(0) &= p\end{aligned}$$

Recall $b^2 = c/m$. We can also think of the IVP as a second order differential equation for $x(t)$.

$$x'' = -b^2x, \quad x(0) = a, \quad x'(0) = 0$$

The Hard Spring

In our derivation of the linear spring model we assumed that the displacement x of the spring was directly proportional to the applied force, to obtain the equation $F = -kx$.

In the non-linear hard spring model, we can have a situation where to in order to double the displacement you have to MORE than double the force, i.e. you have a hard spring, where now the relationship between Force applied and displacement is $F = -cx - \gamma x^3$, $c > 0$, $\gamma > 0$

Sketch the Force versus Displacement graph for the hard spring in the space below:

we can show that the corresponding IVP will be:

$$\begin{aligned}x' &= v & x(0) &= a \\v' &= -b^2x - \beta x^3 & v(0) &= p\end{aligned}$$

What is the behavior of the hard spring like for SMALL displacements, i.e. x much less than 1?

The Pendulum

In the pendulum problem, the relationship between Force and Angular Displacement is $F = -mg \sin(x)$. Sketch the Force versus Displacement graph for the pendulum model in the space below:

The corresponding IVP for the pendulum model will be:

$$\begin{aligned}x' &= v & x(0) &= a \\v' &= -g \sin(x) & v(0) &= p\end{aligned}$$

The pendulum can be considered to behave like a soft spring. Why?

Again, what is the behavior of the pendulum similar to for small values of x ?

If you wanted to graph solutions of these models what method would you use?