

---

# Multivariable Calculus

Math 212 Fall 2005  
©2005 Ron Buckmire

Fowler 307 MWF 9:30pm - 10:25am  
<http://faculty.oxy.edu/ron/math/212/05/>

---

## Class 8: Friday September 23

**SUMMARY** Vector Functions

**CURRENT READING** Williamson & Trotter, Section (Section 4.1 and 4.2)

**HOMEWORK** Williamson & Trotter, page 182: 2, 3, 9, 13, 14, 17

---

---

### DEFINITION

A **vector** function of a **vector** variable  $\vec{f}(\vec{x})$  with **domain**  $D \subset \mathbb{R}^n$  and **range**  $R \subset \mathbb{R}^m$  means that it has possible input values which form a subset of  $\mathbb{R}^n$  and the set of possible output values are a subset of  $\mathbb{R}^m$ . Often the notation  $f : D \rightarrow R$  or  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is used.

In order to produce the output of this vector function of a vector requires  $m$  **coordinate functions** which are scalar functions of a vector variable:  $\vec{f}(\vec{x}) = (f_1(\vec{x}), f_2(\vec{x}), \dots, f_m(\vec{x}))$

### Matrix Representation

If  $f$  is a linear vector function then  $f_i(x_1, x_2, x_3, \dots, x_n) = a_{i1}x_1 + a_{i2}x_2 + \dots + a_{in}x_n$  then it can be replaced by a simple matrix-vector product  $\vec{f}(\vec{x}) = A\vec{x}$

### EXAMPLE 1

The function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by  $y_1(x_1, x_2) = -x_1 - x_2$  and  $y_2(x_1, x_2) = x_1 + 2x_2$  has what matrix form?

### Vector Functions of a Single Variable

Section 4.1 and 4.2 deal with the calculus of vector functions of a single variable which have the form  $f : \mathbb{R} \rightarrow \mathbb{R}^n$ . We've previously seen an example of such a function,  $f : \mathbb{R} \rightarrow \mathbb{R}^3$

### EXAMPLE 2

What kind of geometric object is the image of the function  $\vec{x}(t) = (1 + 3t, -1 - t, -2 + t)$ ?

## Derivative of a Vector Function of a Single Variable

Basically vector functions of a single variable should be treated as collections of real functions of a real variable.

### EXAMPLE 3

Consider  $f : \mathbb{R} \rightarrow \mathbb{R}^4$  where  $\vec{f}(t) = \begin{bmatrix} t^2 + 1 \\ \sin(t) \\ e^{-t} \\ 4 \end{bmatrix}$ . Compute  $\vec{f}(0)$ ,  $\frac{d\vec{f}}{dt}$  and  $\int \vec{f}(t)dt$

### DEFINITION

If a curve has a parametric representation  $\vec{f}(t)$  such that the image of  $\vec{f}$  has a derivative  $\vec{f}'(t)$  is continuous **and** never zero, then the curve is described as **smooth**.

### EXERCISE 1

Show that the curve defined by  $\vec{x}(t) = (\cos(t), \sin(t), t)$  is a **helix** and is a **smooth curve**.

## Tangent Vector and Tangent Line

The tangent vector to a parametric curve  $\vec{x}(t)$  is given by the derivative of  $\vec{x}$  with respect to  $t$  and can be denoted  $\dot{\mathbf{x}}$ . Using this information, it's clear that the equation of a tangent line  $\vec{t}(t)$  to a curve  $\vec{x}(t)$  at the point  $\mathbf{x}_0$  is given by  $\vec{t}(t) = \dot{\mathbf{x}}t + \mathbf{x}_0$

### EXERCISE 2

Find the equation of the tangent line to the helical curve  $\vec{x}(t)$  from the previous example at the point  $\mathbf{x}_0 = (1, 0, 0)$ .

### DEFINITION

Length of a curve  $l(\gamma) = \int_{t_0}^{t_1} |\dot{\mathbf{x}}|dt$  where  $\gamma$  is the parameterized curve between  $t = t_0$  and  $t = t_1$ .

### EXERCISE 3

Find an expression for the length of one revolution of a helix from  $t = 0$  to  $t = 2\pi$ .