

---

# Multivariable Calculus

Math 212 Fall 2005  
©2005 Ron Buckmire

Fowler 307 MWF 9:30pm - 10:25am  
<http://faculty.oxy.edu/ron/math/212/05/>

---

*Class 10: Wednesday September 28*

**SUMMARY** Partial Derivatives

**CURRENT READING** Williamson & Trotter, Section 4.3

**HOMEWORK** Williamson & Trotter, page 203: 3, 9, 12, 22, 25, 31, 34, 37; **Extra Credit page 204: # 38, 39**

---

---

## DEFINITION

The **partial derivative** of a scalar function of a vector variable  $f(\vec{x})$  where  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  can be defined as

$$\frac{\partial f}{\partial x_i} = \lim_{h \rightarrow 0} \frac{f(x_1, x_2, x_i + h, \dots, x_n) - f(x_1, x_2, \dots, x_n)}{h}$$

In practice, this basically means that if you have a function of many variables when you take a partial derivative with respect to a particular variable you treat all the other variables in the function as **CONSTANTS**.

**NOTE:** Sometimes  $\frac{\partial f}{\partial x}$  will be denoted simply  $f_x$ .

## EXAMPLE 1

Consider  $f(x, y, z) = xyz + \sin(x + y) + z^2y + e^{-x}$  and calculate  $f_x$ ,  $f_y$  and  $f_z$

## Exercise 1

Williamson & Trotter, page 204, page 21. Find  $\frac{\partial^3 f}{\partial x^2 \partial y}$  if  $f(x, y) = \ln(x + y)$

### Equation of a Tangent Plane to a Surface

The equation of a tangent plane to a surface  $z = f(x, y)$  at the point  $(a, b, f(a, b))$  is given by the equation

$$z = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

#### Exercise 2

**Williamson & Trotter, page 204, page 10.** Find the equation of the tangent plane to  $f(x, y) = x(y^2 + 1)$  at  $(a, b) = (0, 2)$ .

### Graphical Interpretation of Tangent Plane

The notion of the existence of a tangent plane to a surface is the 3-dimension equivalent to the existence of a tangent line or **local linearity** of a curve in 2-dimensions. The existence of these objects relate to the **differentiability** of the scalar function  $f(\vec{x})$ , and **differentiability of a function at a point implies continuity at that point**, but continuity DOES NOT imply differentiability.

### Connection to Taylor's Theorem and Microscope Approximation

We can approximate a function  $f(x, y)$  near the point  $(x_0, y_0)$  with its tangent plane:

$$f(x, y) \approx f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

#### EXAMPLE 2

Approximate the value of  $f(2.01, 0.97)$  where  $f(x, y) = \sqrt{x^2 + y^3}$ .

#### Clairault's Theorem

If  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  is continuous and  $f_x, f_y, f_{xy}$  and  $f_{yx}$  are also continuous on the same domain as  $f$ , then  $f_{xy} = f_{yx}$ . **NOTE:**  $f_{xy} = (f_x)_y$ .

#### Exercise 3

Show that Clairault's Theorem applies to the function  $f(x, y) = x^y$ .