

---

# Multivariable Calculus

Math 212 Spring 2006  
©2006 Ron Buckmire

Fowler 112 MWF 8:30pm - 9:25am  
<http://faculty.oxy.edu/ron/math/212/06/>

---

*Class 3: Friday January 27*

**SUMMARY** The Dot Product and its Implications and Applications

**CURRENT READING** Williamson & Trotter, Section 1.4 and 1.5

**HOMEWORK #3** Williamson & Trotter, Page 36 # 5, 6, 7, 8, 9, 18, 21, 22, 27;

---

---

## Dot Product

Given two vectors in  $\mathbb{R}^n$ ,  $\vec{x} = (x_1, x_2, \dots, x_n)$  and  $\vec{y} = (y_1, y_2, \dots, y_n)$  the **dot product** is defined as:

$$\vec{x} \cdot \vec{y} = \sum_{k=1}^n x_k y_k = x_1 y_1 + x_2 y_2 + x_3 y_3 + \dots + x_n y_n$$

The dot product is a very useful operation that allows us to represent a number of interesting results.

## Magnitude of a Vector

$$|\vec{x}| = \sqrt{\vec{x} \cdot \vec{x}}$$

## Angles Between Vectors

The dot product also defines an expression for the angle between two vector  $\vec{x}$  and  $\vec{y}$

$$\vec{x} \cdot \vec{y} = |\vec{x}| |\vec{y}| \cos(\theta)$$

which leads to the **Cauchy-Schwarz Inequality**

$$\vec{x} \cdot \vec{y} \leq |\vec{x}| |\vec{y}|$$

## Law of Cosines

$$|\vec{x} - \vec{y}|^2 = |\vec{x}|^2 + |\vec{y}|^2 - 2|\vec{x}| |\vec{y}| \cos(\theta)$$

## Triangle Inequality

$$|\vec{x} + \vec{y}| \leq |\vec{x}| + |\vec{y}|$$

## Properties of the Dot Product

**Positivity:**  $\vec{x} \cdot \vec{x} > 0$  (except when  $\vec{x} = \vec{0}$ )

**Symmetry:**  $\vec{x} \cdot \vec{y} = \vec{y} \cdot \vec{x}$

**Additivity:**  $(\vec{x} + \vec{y}) \cdot \vec{z} = \vec{x} \cdot \vec{z} + \vec{y} \cdot \vec{z}$

**Homogeneity:**  $(r\vec{x}) \cdot \vec{y} = r(\vec{x} \cdot \vec{y})$

**GROUPWORK**

For the given vector  $\vec{u} = (3, 1, 1)$  and  $\vec{v} = (4, 1, 0)$  find  $\vec{u} \cdot \vec{v}$ ,  $|\vec{u}|$ ,  $|\vec{v}|$  the angle between  $\vec{v}$  and  $\vec{u}$  and normalize each of the vectors.

**EXERCISE**

**Williamson & Trotter, page 32, # 28.** Show that the sum of the squares of the lengths of the four sides of a parallelogram is equal to the sum of the squares of the diagonals. [HINT: Sketch the two vectors  $\vec{x}$  and  $\vec{y}$  and obtain expressions for the diagonals in terms of  $\vec{x}$  and  $\vec{y}$ .]