

---

# Linear Systems

Math 214 Spring 2006  
©2006 Ron Buckmire

Fowler 307 MWF 2:30pm - 3:25pm  
<http://faculty.oxy.edu/ron/math/214/06/>

---

*Class 17: Monday March 6*

**TITLE** Subspaces Associated With Matrices; Dimension and Basis

**CURRENT READING** Poole 3.5 and 6.1

## Summary

Let's continue discussing vector spaces associated with matrices and formally define the concept of dimension.

---

---

## Homework Assignment

Poole, Section 3.5: **17, 18, 21, 24, 39, 40, 41, 42**. EXTRA CREDIT 44, 50.

---

---

Recall: Let  $A$  be an  $m \times n$  matrix. The **row space** of  $A$  is the subspace of  $\mathbb{R}^n$  spanned by the rows of  $A$  and is denoted  $\text{row}(A)$ . The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$  and is denoted  $\text{col}(A)$ .

## Warm-Up

Given  $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 4 & 7 \end{bmatrix}$ . Find  $\text{col}(A)$  and  $\text{row}(A)$ .

## DEFINITION

The **null space** of a  $m \times n$  matrix is the subspace of  $\mathbb{R}^n$  consisting of all solutions of the homogeneous linear system  $A\vec{x} = \vec{0}$ . It is denoted by  $\text{null}(A)$ .

## Theorem 3.21

The set  $N$  of all solutions to the homogeneous linear system  $A\vec{x} = \vec{0}$  where  $A$  is a  $m \times n$  matrix is a subspace of  $\mathbb{R}^n$ .

## Exercise

Prove Theorem 3.21 (that the nullspace of a matrix  $A$  is a subspace of  $\mathbb{R}^n$ ).

**DEFINITION**

The **basis** of a subspace  $\mathcal{S}$  of  $\mathbb{R}^n$  is a set of vectors in  $\mathcal{S}$  which is **linear independent** and **spans**  $\mathcal{S}$ . The plural of basis is bases.

**Q:** Are bases for a subspace unique? **A:** Heck, no! (Why not?)

**GROUPWORK**

Write down three examples of bases for  $\mathbb{R}^2$ .

**Theorem 3.23**

The number of vectors found in a basis for a subspace  $\mathcal{S}$  of  $\mathbb{R}^n$  is the same. Any two bases for  $\mathcal{S}$  have the same number of vectors.

**DEFINITION**

The number of vectors in a basis for a subspace  $\mathcal{S}$  of  $\mathbb{R}^n$  is known as the **dimension** of  $\mathcal{S}$  and is denoted  $\dim(\mathcal{S})$ . This result is known as the **Basis Theorem**.

**EXAMPLE**

$$\text{Given } A = \begin{bmatrix} 1 & 1 & 3 & 1 & 6 \\ 2 & -1 & 0 & 1 & -1 \\ -3 & 2 & 1 & -2 & 1 \\ 4 & 1 & 6 & 1 & 3 \end{bmatrix} \text{ and } \text{rref}(A) = \begin{bmatrix} 1 & 0 & 1 & 0 & -1 \\ 0 & 1 & 2 & 0 & 3 \\ 0 & 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Let's write down bases for  $\text{col}(A)$ ,  $\text{row}(A)$  and  $\text{null}(A)$ .

**Theorem 3.20**

If  $B$  is a matrix formed from applying elementary row operations to  $A$  and thus  $B$  is row equivalent to  $A$ , then  $\text{row}(A) = \text{row}(B)$ .

Elementary row operations **do not affect** the row space of a matrix, but they **do change** the column space of a matrix. Given  $R = \text{rref}(A)$ .  $\text{row}(A) = \text{row}(R)$  but  $\text{col}(A) \neq \text{col}(R)$ .



**Theorem 3.27**

**The Fundamental Theorem of Invertible Matrices (Version 2).** Let  $A$  be a  $n \times n$  matrix. Each of the following statements is equivalent:

- (a)  $A$  is invertible.
- (b)  $A\vec{x} = \vec{b}$  has a unique solution for every  $\vec{b}$  in  $\mathbb{R}^n$ .
- (c)  $A\vec{x} = \vec{0}$  has only the trivial solution.
- (d) The reduced row echelon form of  $A$ ,  $\text{rref}(A)$ , is  $I_n$ .
- (e)  $A$  is a product of elementary matrices.
- (f)  $\text{rank}(A) = n$ .
- (g)  $\text{nullity}(A) = 0$ .
- (h) The column vectors of  $A$  are linearly independent.
- (i) The column vectors of  $A$  span  $\mathbb{R}^n$ .
- (j) The column vectors of  $A$  form a basis for  $\mathbb{R}^n$ .
- (k) The row vectors of  $A$  are linearly independent.
- (l) The row vectors of  $A$  span  $\mathbb{R}^n$ .
- (m) The row vectors of  $A$  form a basis for  $\mathbb{R}^n$ .

**Standard Basis and Coordinates**

The standard unit vectors in  $\mathbb{R}^n$  are the  $n$  rows and columns of the identity matrix  $I_n$ . A **standard basis** for  $\mathbb{R}^n$  would be a collection of  $n$  of these vectors, usually denoted  $\hat{e}_1, \hat{e}_2, \hat{e}_3, \dots, \hat{e}_n$ .

**Theorem 3.28**

Let  $\mathcal{S}$  be a subspace of  $\mathbb{R}^n$  and let  $\beta = \{\vec{v}_1, \vec{v}_2, \vec{v}_3, \dots, \vec{v}_k\}$  be a basis for  $\mathcal{S}$ . For every vector  $\vec{v}$  in  $\mathcal{S}$  there is exactly one way to write  $\vec{v}$  as a linear combination of the basis vectors in  $\beta$ :

$$\vec{v} = c_1\vec{v}_1 + c_2\vec{v}_2 + c_3\vec{v}_3 + \dots + c_k\vec{v}_k$$

**DEFINITION**

These numbers  $c_1, c_2, \dots, c_k$  are called the **coordinates of  $\vec{v}$  with respect to  $\beta$** .

The vector  $[\vec{v}]_\beta = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_k \end{bmatrix}$  is known as the **coordinate vector of  $\vec{v}$  with respect to  $\beta$** .

**EXAMPLE**

**Poole, page 209, #49.** Show that  $(1, 6, 2)$  is in  $\text{span}(\beta)$ , where  $\beta = \{(1, 2, 0), (1, 0, -1)\}$  and find the coordinate vector  $[\vec{w}]_\beta$ .