Multivariable Calculus

Math 224 Spring 2004 ©2004 Ron Buckmire

Fowler 112 MWF 2:30pm - 3:25pm http://faculty.oxy.edu/ron/math/224/04/

Class 19: Monday March 8

SUMMARY Implicit Differentiation CURRENT READING Williamson & Trotter, Section 6.3 HOMEWORK #18 Williamson & Trotter, page 274: 2, 3; page 281: 2, 3, 4, 5, 7, 12, 15

THEOREM: The Inverse Function Theorem

Let $\vec{F} : \mathbb{R}^n \to \mathbb{R}^n$ be continuously differentiable on an open subset S of \mathbb{R}^n and let \vec{x}_0 be a point in S with invertible derivative matrix (Jacobian) $\vec{F'}$. THEN there is a neighborhood N of \vec{x}_0 such that \vec{F} has a continuously differentiable inverse function $(\vec{F})^{-1}$ defined on the image set $\vec{F}(N)$. The derivative matrix (Jacobian) of \vec{F}^{-1} is related to the Jacobian matrix of \vec{F} by the equation

$$[\vec{F}^{-1}]'(\vec{F}(\vec{x})) = [\vec{F}'(\vec{x})]^{-1}$$

You should read this as the derivative of the inverse function of \vec{F} evaluated at \vec{F} is equal to the inverse of the derivative of the function \vec{F} evaluated at \vec{x} .

This is the multivariable version of the common result from Calculus you may recall that given f(a) = b and a = g(b) so that g is the inverse function of f, then $g'(f(a)) = g'(b) = \frac{1}{f'(a)}$ <u>EXAMPLE 1</u> $f(x) = e^x$ and $g(x) = \ln(x)$ then f and g are inverse functions of each other. Let a = 1 and $b = f(a) = e^1 = e$. Compute g'(e) using the Inverse Function Theorem.

The main usefulness of the Inverse Function Theorem is when one is doing a **coordinate transformation**. This will be extremely more important when we look at Multivaiable Integration later on.

EXAMPLE 2

Williamson & Trotter, page 274, #10. Define $P : \mathbb{R}^2 \to \mathbb{R}^2$ by the equations $x = u \cos v, y = \sin v$ for u > 0.

(a) Show that for fixed $v = v_0$ and varying u > 0 the image curves in the xy-plane are half-lines emanating from x, y = (0, 0).

(b) Show that for fixed $u = u_0$ and varying v the image curves in the xy-plane are circles of radius u_0 each one traced infinitely often.

(c) Compute the determinant of the jacobian matrix P'(u, v) and show that if $u_0 \neq 0$ then inverse function theorem implies the existence of a local inverse in the neighborhood of $(x_0, y_0) = P(u_0, v_0)$.

Implicit Differentiation

Recall that even though one does not always have an explicit definition of a curve y = f(x) one can still compute the slope of such a curve using a process called **implicit differentia**tion. The curve is defined *implicitly* as F(x, y) = c.

Using the Chain Rule this equation becomes $\frac{\partial F}{\partial x}\frac{dx}{dx} + \frac{\partial F}{\partial y}\frac{dy}{dx} = 0$ when when solved implies

that
$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -(F_y)^{-1}F_x$$

Exercise 1

The equation $F(x,y) = x^2 + y^2 - 4 = 0$ defines a circle of of radius 2 centered at the origin. What's the slope of this curve at $(1,\sqrt{3})$? What about at (2,2)?

THEOREM: The Implicit Function Theorem

Let $F : \mathbb{R}^{n+m} \to \mathbb{R}^m$ be a continuously differentiable function. Suppose for some \vec{x}_0 in \mathbb{R}^n and some \vec{y}_0 in \mathbb{R}^m that

(i) $\vec{F}(\vec{x}_0, \vec{y}_0) = \vec{0}$ and

(ii) $\vec{F}_{\vec{y}}(\vec{x}_0, \vec{y}_0)$ is an *m*-by-*m* invertible matrix.

THEN there is a unique continuously differentiable function $G : \mathbb{R}^n \to \mathbb{R}^m$ defined on an open neighborhood N of \vec{x}_0 in \mathbb{R}^n such that $\vec{F}(\vec{x}, \vec{G}(\vec{x})) = \vec{0}$ for all \vec{x} in N and $\vec{G}(\vec{x}_0) = \vec{y}_0$.

Theorem

Suppose $F : \mathbb{R}^{n+m} \to \mathbb{R}^m$ and $G : \mathbb{R}^n \to \mathbb{R}^m$ are differentiable and that $\vec{y} = \vec{G}(\vec{x})$ satisfies $\vec{F}(\vec{x}, \vec{y}) = \vec{0}$ for all \vec{x} in some open subset of \mathbb{R}^n . THEN

$$\vec{G}'(\vec{x}) = -[\vec{F}_{\vec{y}}]^{-1}(\vec{x}, \vec{G}(\vec{x}))\vec{F}_{\vec{x}}(\vec{x}, \vec{G}(\vec{x}))$$

Exercise 2

Williamson & Trotter, page 281, #8. If x + y + u - v = 0 and x + y + 2u + v = 0. Find $\frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial x}{\partial v}$ and $\frac{\partial y}{\partial v}$ by (1) solving for x and y in terms of u and v and (2) by implicit differentiation.

EXAMPLE 3

Consider Williamson & Trotter, page 281, #9. Suppose $x^2y + yz = 0$ and xyz + 1 = 0. Compute $\vec{F}, \vec{x}, \vec{y}$ and $\vec{F}_{\vec{x}}$ and $\vec{F}_{\vec{y}}$ at (1, 1, -1).