Real Analysis

Math 310 — MWF 10:30-11:25am
Fall 1996 — 127 Fowler

Ron Buckmire ~ 402 Fowler ~ 259-2536 ~ ron@oxy.edu ~ Mon 1:30-3:30pm Wed & Thu 1:30-2:30pm

---

#27 Friday (11/22/96)

SUMMARY Uniform Continuity

READING §19 (page 102-110)

Definition
A function $\tilde{f}$ is said to be an **extension of** f if $\text{dom}(f) \subseteq \text{dom}(\tilde{f})$ and $f(x) = \tilde{f}(x)$ for all $x \in \text{dom}(f)$
If $f(x) = \sin(1/x)$ for $x \in (0, 1/\pi]$ Observe this extension of f written below:

$$\tilde{f}(x) = \begin{cases} x\sin(1/x) & \text{for} \quad 0 < x \leq 1/\pi \\ 0 & \text{for} \quad x = 0 \end{cases}$$

What is $\text{dom}(f)$?
What is $\text{dom}(\tilde{f})$?
Is $\tilde{f}$ continuous at $x = 0$?

Theorem
19.5 A real-valued function on (a,b) is uniformly continuous on (a,b) if and only if it can be extended to a continuous function $\tilde{f}$ on [a,b]

Theorem
19.6 Let f is a continuous function on an interval I (I may be bounded or unbounded). Let I^0 is the interval unbounded by removing any endpoints of I which happen to be in I. If f is **differentiable** on I^0 and if f' is **bounded on** I^0, then f is **uniformly continuous** on I.
PROOF: Let's read thru page 108 of the text.

Homework Assignment (Due Friday November 22)
§19, page 109: **19.1, 19.2, 19.4(a), 19.5(d),(e),(f)**