

---

# Complex Analysis

Math 214 Spring 2004  
©2004 Ron Buckmire

Fowler 112 MWF 3:30pm - 4:25pm  
<http://faculty.oxy.edu/ron/math/312/04/>

---

## *Class 7: Wednesday February 4*

**SUMMARY** Limits of Complex Functions and the Point At Infinity

**CURRENT READING** Saff & Snider, §2.2

**HOMEWORK** Saff & Snider, Section 2.2 # 1, 5, 7, 11, 14 **Extra Credit: #25**

---

---

### **Limits**

Suppose that  $f(z)$  is defined on a deleted neighborhood of  $z_0$ . In order to say that  $\lim_{z \rightarrow z_0} f(z) = w_0$  we must be able to show that

$$\forall \epsilon > 0, \quad \exists \delta > 0 \ni |z - z_0| < \delta \Rightarrow |f(z) - w_0| < \epsilon$$

This may look like dense mathematical language, but in english this means that for every positive number  $\epsilon$  (no matter how small) there exists a number  $\delta$  (which depends on the choice of  $\epsilon$ ) so that regardless of how close you get to the point  $z_0$  in the deleted neighborhood around it in the  $z$ -plane you can also get arbitrarily close to the value  $w_0$  in the  $w$ -plane.

### **Visualize The Limit**

Let's try and prove the result  $\lim_{z \rightarrow i} z^2 = -1$  (See Example 2 on page 60 of Saff & Snider)

## Rules on Limits

The rules on limits of complex functions are identical to the rules for limits of real functions of real variables (as you'd expect)

Suppose that  $\lim_{z \rightarrow z_0} f(z) = w_0$  and  $\lim_{z \rightarrow z_0} F(z) = W_0$  then

$$\lim_{z \rightarrow z_0} [f(z) + F(z)] = w_0 + W_0$$

$$\lim_{z \rightarrow z_0} [f(z)F(z)] = w_0W_0$$

$$\lim_{z \rightarrow z_0} \frac{f(z)}{F(z)} = \frac{w_0}{W_0} \quad (W_0 \neq 0)$$

$$\lim_{z \rightarrow z_0} |f(z)| = |w_0|$$

$$\lim_{z \rightarrow z_0} c = c$$

$$\lim_{z \rightarrow z_0} z^n = z_0^n$$

$$\lim_{z \rightarrow z_0} P(z) = P(z_0) \quad (\text{where } P(z) \text{ is a polynomial})$$

IF  $f(z) = u(x, y) + iv(x, y)$ ,  $z_0 = x_0 + iy_0$  and  $w_0 = u_0 + iv_0$  THEN  
 $\lim_{z \rightarrow z_0} f(z) = w_0$  if and only if  $\lim_{(x,y) \rightarrow (x_0,y_0)} u(x, y) = u_0$  and  $\lim_{(x,y) \rightarrow (x_0,y_0)} v(x, y) = v_0$

## Examples

(a)  $\lim_{z \rightarrow 1+2i} 2|z| + iz^2 + 2.5 - .1i =$

(b)  $\lim_{z \rightarrow 3\pi i} ze^z =$

(c)  $\lim_{z \rightarrow 0} \frac{z^8 + z^4 + z^2 + z - 1}{z^3 + 4z^3 - 9} =$

(d)  $\lim_{z \rightarrow 1-i} 2xy - ix^2 - iy^2 =$

## Point at Infinity

When dealing with real numbers we often speak of two different concepts, denoted  $-\infty$  and  $+\infty$ .

However, the complex number infinity is represented as one particular point in the Argand plane. We call this the **point of infinity** and we rename the Argand plane the **extended  $z$  plane** or the **extended complex plane** when we include it.

The point at infinity can be considered to be the image of the origin  $z = 0$  under the mapping  $w = 1/z$

To compute complex limits involving  $\infty$  we use this idea:

$$\lim_{z \rightarrow \infty} f(z) = \lim_{w \rightarrow 0} f\left(\frac{1}{w}\right)$$

In order for us to say a complex function  $f(z)$  becomes unbounded at a point  $z_0$ , (i.e.  $f(z_0) = \infty$ ) we must be able to show that

$$\forall M > 0, \quad \exists \delta > 0 \quad \exists 0 < |z - z_0| < \delta \Rightarrow |f(z)| > M$$

## Examples

(a)  $\lim_{z \rightarrow \infty} \frac{2z + 3}{3z - i} =$

(b)  $\lim_{z \rightarrow \infty} z^2 + 3z - 5i + 4 =$