
Complex Analysis

Math 312 Spring 1998
Buckmire

MWF 10:30am - 11:25am
Fowler 112

Class 10 (Wednesday February 4)

SUMMARY Analyticity and the Cauchy-Riemann Equations

CURRENT READING Brown & Churchill, pages 45-57

NEXT READING Brown & Churchill, pages 48-50, 55-57, 59-64

GROUPWORK

Given $g(z) = z^2 + z + i$ and $f(z) = \frac{1}{z}$

$g'(z) =$

$f'(z) =$

$[g(z)f(z)]' =$

$[g(z)/f(z)]' =$

Cauchy-Riemann Equations

We shall derive the Cauchy-Riemann equations by looking at the definition of the derivative of a function $f(z) = u(x, y) + iv(x, y)$ at the point z_0 .

$$\begin{aligned} f'(z_0) &= \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} \\ &= \lim_{(\Delta x, \Delta y) \rightarrow (0,0)} \frac{u(x_0 + \Delta x, y_0 + \Delta y) + iv(x_0 + \Delta x, y_0 + \Delta y) - u(x_0, y_0) - iv(x_0, y_0)}{\Delta x + i\Delta y} \end{aligned}$$

We shall do this limit twice, once letting $\Delta z \rightarrow 0$ horizontally and the other time letting $\Delta z \rightarrow 0$ vertically

This shows us that the existence of $f'(z_0)$ implies the Cauchy-Riemann equations are satisfied at this point (and at every point in a neighborhood of z_0).

This is true since if $f'(z_0)$ exists then f is **analytic** "at" this point.

ANALYTICITY \Rightarrow C.R.E.

To make satisfying the CRE a *sufficient* condition one needs the added condition that the first derivatives of u and v are continuous. If both these conditions are true and f is defined on an open set, then f is analytic on the open set.

$$f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0) = -i(u_y(x_0, y_0) + iv_y(x_0, y_0))$$

ANALYTICITY \iff C.R.E. + Continuity of u_x, u_y, v_x, v_y

Example

Show that $f(z) = \bar{z}$ is **not analytic** anywhere in the complex plane. You can do this in two ways:

1:

2:

GROUPWORK

Example

Show that the function $f(z) = 1/z$ is analytic on the set $z \neq 0$. To do that you will have to answer the following questions:

- What is its domain of definition? Is this an open set?
- What are its component functions? Are there partial derivatives continuous?
- Do they satisfy the CRE?
- Is it analytic? On what set?