

Name: _____

Monday November 24
Prof. Ron Buckmire

Topic : Convolution!

The idea behind this quiz is to provide you with an opportunity to illustrate your understanding of solution curves of linear systems in 2-D.

Reality Check:

EXPECTED SCORE : _____/5

ACTUAL SCORE : _____/5

Instructions:

0. Please look for a hint on this quiz posted to faculty.oxy.edu/ron/math/341/08/
1. Once you open the quiz, you have **30 minutes** to complete, please record your start time and end time at the top of this sheet.
2. You may use the book or any of your class notes. You must work alone. **Do not mention the existence of this quiz to anyone else in the class.**
3. If you use your own paper, please staple it to the quiz before coming to class. If you don't have a stapler, buy one. QUIZZES WITH UNSTAPLED SHEETS WILL NOT BE GRADED.
4. After completing the quiz, sign the pledge below stating on your honor that you have adhered to these rules.
5. Your solutions must have enough details such that an impartial observer can read your work and determine HOW you came up with your solution.
6. Relax and enjoy...
7. **This bonus quiz is due on Monday December 1**, with the rest of your week's homework. **NO LATE OR UNSTAPLED QUIZZES WILL BE ACCEPTED.**

Pledge: I, _____, pledge my honor as a human being and Occidental student, that I have followed all the rules above to the letter and in spirit.

1. Consider the following “auto-convolution” equation

$$f * f = \int_0^t f(\tau)f(t - \tau) d\tau = 6t^3$$

(a) *5 points.* **Show** that the function $f(t) = \pm 6t$ is the solution of the above equation. (Solve the integral equation using Laplace Transforms and you should find that the solution is indeed the given function.)

(b) *5 points.* **Confirm** that if $f(t) = \pm 6t$, then the convolution of $f(t)$ with itself is $6t^3$. (Check that the given function indeed satisfies the given equation when “auto-convolved.”)