

---

# Differential Equations

Math 341 Fall 2014  
©2014 Ron Buckmire

MWF 3:00-3:55pm Fowler 307  
<http://faculty.oxy.edu/ron/math/341/14/>

---

## Worksheet 25

**TITLE** Introducing The Laplace Transform

**CURRENT READING** Blanchard, 6.1

---

---

**Homework #11 Assignments due Monday November 17**

**Section 6.1:** 2, 3, 5, 7, 8, 9, 15, 18, 25\*.

**Section 6.2:** 1, 2, 4, 8, 15, 16, 18\*.

**Homework #12 Assignments due Monday November 24**

Section 6.3: 5, 6, 8, 15, 18, 27, 28.

Section 6.4: 1, 2, 6, 7\*.

---

---

### SUMMARY

We introduce a new kind of operator, an integral operator, called the Laplace Transform, which can be used to solve differential equations.

---

## 1. Introducing The Laplace Transform

### DEFINITION: Integral Transform

If a function  $f(t)$  is defined on  $[0, \infty)$  then we can define an integral transform to be the improper integral  $F(s) = \int_0^{\infty} K(s, t)f(t) dt$ . If the improper integral converges then we say that  $F(s)$  is the integral transform of  $f(t)$ . The function  $K(s, t)$  is called the **kernel** of the transform. When  $K(s, t) = e^{-st}$  the transform is called **the Laplace Transform**.

### DEFINITION: Laplace Transform

Let  $f(t)$  be a function defined on  $t \geq 0$ . The Laplace Transform of  $f(t)$  is defined as

$$F(s) = \mathcal{L}[f(t)] = \int_0^{\infty} e^{-st} f(t) dt$$

(Note the use of capital letters for the transformed function and the lower-case letter for the input function.) Some people use curly brackets to denote the input, like  $\mathcal{L}\{f(t)\}$  but we will use the textbook's notation of square bracket.)

**EXAMPLE** Let's show that  $\mathcal{L}[1] = \frac{1}{s}, s > 0$

### Exercise

Compute  $\mathcal{L}[t]$ .

## 2. Linearity Property of The Laplace Transform

$\mathcal{L}$  is a linear operator, in other words  $\mathcal{L}[af(t) + bg(t)] = a\mathcal{L}[f(t)] + b\mathcal{L}[g(t)]$

**EXAMPLE** Let's prove the Laplace Transform possesses the linearity property.

**Q:** Does every function have a Laplace Transform?

**A:** Hell, no! (i.e.  $t^{-1}$ ,  $e^{t^2}$  etc do not). Can you think of any others?

**DEFINITION: exponential order**

A function  $f$  is said to be of **exponential order**  $c$  if there exist constants  $c$ ,  $M > 0$ ,  $T > 0$  such that  $|f(t)| \leq Me^{ct}$  for all  $t > T$ .

Basically this is saying that in order for  $f(t)$  to have a Laplace Transform then in a race between  $|f(t)|$  and  $e^{ct}$  as  $t \rightarrow \infty$  then  $e^{ct}$  must approach its limit first, i.e.  $\lim_{t \rightarrow \infty} \frac{f(t)}{e^{ct}} = 0$ .

**THEOREM**

If  $f$  is piecewise continuous on  $[0, \infty)$  and of exponential order  $c$ , then  $F(s) = \mathcal{L}[f(t)]$  exists for  $s > c$  and  $\lim_{s \rightarrow \infty} F(s) = 0$

This result means that there are functions that clearly can NOT BE Laplace Transforms. These would be functions who do not satisfy the conclusion of the above theorem.

**GROUPWORK**

Which of the following functions can NOT be Laplace Transforms? Which of the following MIGHT be Laplace Transforms?(HINT: think of the contrapositive of the theorem!)

(a)  $\frac{s}{s+1}$

(b)  $\frac{s}{s^2+1}$

(c)  $\frac{s^2}{s+1}$

(d)  $s^2 + 1$

### 3. Laplace Transforms of Piecewise Continuous Functions

**Exercise** Find the Laplace Transform of the piecewise function  $f(t) = \begin{cases} 0, & 0 \leq t < 3 \\ 2, & t \geq 3 \end{cases}$

### 4. Transforming A Derivative

**EXAMPLE** We can show that  $\mathcal{L}[f'(t)] = sF(s) - f(0)$

**THEOREM**

If  $f, f', f'', f^{(n-1)}, \dots, f^{(n-1)}$  are continuous on  $[0, \infty)$  and of exponential order  $c$  and if  $f^{(n)}$  is piecewise continuous on  $[0, \infty)$ , then  $\mathcal{L}[f^{(n)}(t)] = s^n F(s) - \sum_{k=1}^n s^{n-k} f^{(k-1)}(0)$

### 5. Using Transforms To Solve Differential Equations

**EXAMPLE** Blanchard, Devaney & Hall, page 577, #15. Use the Laplace Transform to solve the initial value problem  $\frac{dy}{dt} = -y + e^{-2t}, \quad y(0) = 2$

## 6. The Inverse Laplace Transform

### DEFINITION: Inverse Laplace Transform

If  $F(s)$  represents the Laplace Transform of a function  $f(t)$  such that  $\mathcal{L}[f(t)] = F(s)$  then the Inverse Laplace Transform of  $F(s)$  is  $f(t)$ , i.e.  $\mathcal{L}^{-1}[F(s)] = f(t)$ .

Laplace Transforms		Inverse Laplace Transforms	
$f(t)$	$F(s) = \mathcal{L}[f(t)]$	$F(s)$	$f(t) = \mathcal{L}^{-1}[F(s)]$
1	$\frac{1}{s}$	$\frac{1}{s}$	1
$t^n$	$\frac{n!}{s^{n+1}}$	$\frac{1}{s^{n+1}}$	$\frac{t^n}{n!}$
$e^{at}$	$\frac{1}{s-a}$	$\frac{1}{s-a}$	$e^{at}$
$\sin(kt)$	$\frac{k}{s^2+k^2}$	$\frac{k}{s^2+k^2}$	$\sin(kt)$
$\cos(kt)$	$\frac{s}{s^2+k^2}$	$\frac{s}{s^2+k^2}$	$\cos(kt)$
$\sinh(kt)$	$\frac{k}{s^2-k^2}$	$\frac{k}{s^2-k^2}$	$\sinh(kt)$
$\cosh(kt)$	$\frac{s}{s^2-k^2}$	$\frac{s}{s^2-k^2}$	$\cosh(kt)$
$\frac{dg}{dt}$	$sG(s) - g(0)$	$sG(s) - g(0)$	$\frac{dg}{dt}$

### Exercise

Compute  $\mathcal{L}^{-1}\left[\frac{1}{s^5}\right]$  and  $\mathcal{L}^{-1}\left[\frac{1}{s^2+7}\right]$

### EXAMPLE

Let's show that  $\mathcal{L}^{-1}\left[\frac{-2s+6}{s^2+4}\right] = -2\cos(2t) + 3\sin(2t)$